

5 PREPARATION AND COMPOSITION OF INTER-ALPHA INHIBITOR PROTEINS FROM
HUMAN PLASMA FOR THERAPEUTIC USE

RELATED APPLICATION

This application contains subject matter that is related to that disclosed in provisional
10 patent application Ser. No. 60/_____ filed November 8, 2003, entitled, "Preparation and
Composition of Inter-alpha Inhibitor Proteins from Human Plasma for Therapeutic Use," the
disclosure of which application is incorporated herein in its entirety by this reference.

GOVERNMENT SUPPORT

15 A portion of this invention may have been supported by National Institutes of Health
grants RO1 GM053008, R01 GM057468, and R43 GM065667.

BACKGROUND OF THE INVENTION

The inter-alpha inhibitor protein (I α I ρ) family is a group of plasma-associated serine
20 protease inhibitors. Members of this family are composed of heavy and light polypeptide
subunits that are covalently linked by a glycosaminoglycan. The light chain, also called bikunin,
is responsible for the serine protease inhibitory activity of the molecules. The name "bikunin"
reflects the presence of 2 protease-inhibiting domains of the Kunitz type. In normal plasma,
bikunin is found mostly in a complex form as inter-alpha inhibitor (I α I), which has a molecular
25 weight of 225 kDa, and pre-alpha inhibitor (P α I), which has molecular weight of 120 kDa. In
I α I, bikunin is linked to 2 heavy polypeptide chains, H1 and H2, whereas, in P α I, only a single
heavy chain (H3) is linked to bikunin. In these complexed forms, bikunin remains inactive until
its release by partial proteolytic degradation, a mechanism that serves as a means to regulate
activity. After cleavage from the complex, the activated bikunin is cleared rapidly from plasma
30 by glomerular filtration, a process that is facilitated by its low molecular weight and by receptor-
mediated uptake. US Patent Nos. 6,489,128 and 6,660,482 are related to the use of diagnosing
cancer and sepsis, respectively. Methods of inhibiting metastases and of treating sepsis are also

disclosed, however, the compositions were not substantially pure and had stability, i.e., short half-lives, problems.

Despite the introduction of antibiotics over fifty years ago, which indeed saw a decline of sepsis-induced mortality from 55% to 35%, medicine has not benefited from a significant
5 reduction in mortality of subjects with sepsis. In fact, sepsis continues to be one of the leading causes of death in intensive care units and a large number of septic subjects die of ensuing septic shock and multiple organ failure. Sepsis is a systemic response to infection, e.g., a bacterial infection. It is commonly caused by endotoxins from Gram negative bacteria or exotoxins from Gram positive bacteria (which can trigger endotoxin-like responses). The systemic response can
10 lead to septic shock, which is characterized by a precipitous drop in blood pressure, cardiovascular collapse, and/or multiple organ failure. The mortality rate among subjects diagnosed with septic shock can be as high as 35-45%. Rapidly and reliably treating sepsis has been difficult using conventional medications.

Sepsis and septic shock are associated with activation of innate immunity and coagulation
15 systems. Sepsis and septic shock are characterized clinically by systemic inflammation, coagulopathy, hypotension and multiple organ dysfunction (J.-L. Vincent et al., *Annals of Medicine* 34 (2002) 606-613). During severe sepsis, a network of specific proteases activates clotting, fibrinolytic and complement factors. These proteases can also trigger tissue and organ damage and enhance non-specific proteolysis of clotting and complement factors in plasma (J.
20 Wite et al., *Intensive Care Medicine* 8 (1982) 215-222; S.J. Weiss, *New England Journal of Medicine* 320 (1989) 365-376).

BRIEF SUMMARY OF THE INVENTION

The present invention relates to a process for the purification of inter-alpha inhibitor
25 proteins (α_2 Ip) from human plasma and their use for treatment of human diseases such as sepsis, acute inflammatory diseases, severe shock, septic shock, rheumatoid arthritis, cancer, cancer metastasis, infectious diseases, and preterm labor; or reducing the risk of mortality associated

with sepsis, acute inflammatory diseases, severe shock, septic shock, rheumatoid arthritis, cancer, cancer metastasis, infectious diseases, and preterm labor.

According to one aspect, a process for producing a blood plasma-derived IαIp composition, wherein IαI and PaI are present in the mixture in a physiological proportion
5 comprises isolating from blood plasma a plasma fraction containing IαI and PaI, wherein the IαI and PaI are present in a physiological proportion; and purifying the plasma fraction to obtain an IαIp composition with a purity of IαIp ranging from about 85% to about 100% pure.

According to another aspect a composition of IαIp comprises a mixture of inter-alpha inhibitor protein (IαI) and pre-alpha protein (PaI), wherein the IαI and the PaI are present in said
10 mixture in a physiological proportion ranging from about 85% to about 100% pure.

In a related aspect, a composition of IαIp comprises a mixture of inter-alpha inhibitor protein (IαI) and pre-alpha protein (PaI), wherein the IαI and the PaI are present in said mixture in a physiological proportion and having a high trypsin inhibitory specific activity.

In another related aspect, a composition of IαIp comprises IαI and PaI having a half-life
15 of greater than one hour.

In yet another related aspect, a composition of IαIp comprises IαI and PaI wherein the IαI and PaI are composed of a light chain of inter-alpha inhibitor protein associated with at least one of three heavy chains H1, H2 and H3.

In another related aspect, a composition of IαIp comprises a mixture of inter-alpha
20 inhibitor protein (IαI) and pre-alpha protein (PaI), wherein the IαI and the PaI are present in the mixture in a physiological proportion comprising a light chain of inter-alpha inhibitor protein associated with at least one of four heavy chains H1, H2, H3 and H4.

In still another related aspect, a composition of IαIp is made according to the process, which comprises isolating from blood plasma a plasma fraction containing IαI and PaI, wherein
25 the IαI and PaI are present in a physiological proportion; and purifying the plasma fraction to obtain an IαIp composition with a purity of IαIp ranging from about 85% to about 100% pure.

In another aspect, a pharmaceutical composition according to the invention comprises a therapeutically effective composition of IaIp as described herein and a pharmaceutically acceptable carrier.

Another aspect relates to a method of treating an inflammation related disorder, cancer, or an infectious disease in a subject, which comprises administering a therapeutically effective amount of IaIp produced by any of the processes described *infra*.

In another related aspect, a method of treating a subject comprises determining the pre-treatment level of one or more of IaI, PaI, IaIp, H3, H4, H1, H2, and LC; and administering a therapeutically effective amount of IaIp to the subject.

10 In a related aspect, a method for predicting a response to an IaIp therapy is described. The method comprises assaying a sample obtained from a subject to detect the level of one or more of IaI, PaI, IaIp, H3, H4, H1, H2, and LC; wherein the detected levels identifies a subject that may respond favorable to IaIp therapy.

15 In another related aspect, a method of monitoring the progress of a subject being treated with an IaIp therapy is described and comprises determining the pre-treatment level of one or more of IaI, PaI, IaIp, H3, H4, H1, H2, and LC; administering a therapeutically effective amount of IaIp to the subject; and determining the level of one or more of the levels in the subject after an initial period of treatment with the IaIp, wherein an increase of the level in the subject following treatment with IaIp indicates that the subject is likely to have a favorable clinical response to treatment with IaIp.

20 In another aspect, a kit for IaIp therapy is described and comprises one or more of IaI, PaI, IaIp, H3, H4, H1, H2, and LC; and instructions for therapeutic use.

In a related aspect, a composition is described, which comprises a container including IaIp and a label or package insert with instructions for administering the IaIp to a subject.

25 In a further related aspect, a kit is described, which comprises a composition as described above as well as instructions for therapeutic use.

Other embodiments of the invention are disclosed *infra*.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts histopathology of the spleen. Penicillary artery surrounded by preserved white pulp with extensive loss of red pulp cellular elements in the control animal (A) versus in the
5 IaIp-treated animal (B) with normal surrounding white pulp and slightly hypercellular red pulp (H&E stain, magnification 20x).

Figure 2 depicts the amino acid sequence of H4.

Figure 3 depicts the alterations in bikunin mRNA expression in the liver at 5 and 20 h after cecal ligation and puncture (CLP) or sham operation (Sham). Data are expressed as means
10 \pm SE (n=8/group) and compared by one-way analysis of variance (ANOVA) and Tukey's test: * $P < 0.05$ versus shams. Figure 3a shows the mRNA amplified and separated by size on a gel, and Figure 3b graphically depicts the results with the bikunin levels being normalized to the G3PDH expression.

Figure 4 depicts the alterations in the $t_{1/2}$ of ^{125}I -IaI at 5 and 20 h after cecal ligation and
15 puncture (CLP) or sham operation (Sham). Data are expressed as means \pm SE (n=5/group) and compared by one-way analysis of variance (ANOVA) and Tukey's test:

Figure 5 depicts the alterations in the survival rate at 10 days after cecal ligation and puncture and cecal excision with vehicle treatment (CLP+Vehicle) and cecal ligation and puncture with inter- α -inhibitor treatment (CLP+IaIp). There were 12 animals in each group.
20 The survival rate was estimated by the Kaplan-Meier method and compared by using the log-rank test. * $P < 0.05$ vs. CLP+Vehicle.

Figure 6 depicts the alterations in the survival rate at 10 days after cecal ligation and puncture and cecal excision with vehicle treatment (CLP+Vehicle) and cecal ligation and puncture with inter- α -inhibitor treatment (CLP+IaIp). There were 11 to 12 animals in each
25 group. The survival rate was estimated by the Kaplan-Meier method and compared by using the log-rank test. * $P < 0.05$ vs. CLP+Vehicle.

Figure 7 describes alterations in the survival rate at 10 days after cecal ligation and puncture and cecal excision with vehicle treatment (CLP+Vehicle) and cecal ligation and puncture with inter- α -inhibitor treatment (CLP+I α Ip). There were 16 animals in each group. The survival rate was estimated by the Kaplan-Meier method and compared by using the log-rank test. * P < 0.05 vs. CLP+Vehicle.

Figures 8a and 8b describe purification schemes of I α Ip from human plasma, according to the invention.

Figure 9 graphically depicts the beneficial effects of highly purified I α Ip in the animal studies of sepsis.

DETAILED DESCRIPTION OF THE INVENTION

Disclosed herein is a new process for purifying I α Ip from plasma. Further disclosed herein is a therapeutic composition of purified I α Ip for administration to a subject to treat acute inflammatory disease, sepsis, severe shock, septic shock, rheumatoid arthritis, cancer, cancer metastasis, infectious disease, and preterm labor.

Prior known purifications contained contamination by factor X (FX), which is detected as an 80 kDa band by Western blot analysis. The FX was also detected in a clotting assay. The removal of FX is important because FX is considered thrombogenic and can be harmful if administered to humans. The methods described herein solve the FX contamination problems of previous I α Ip compositions.

Inter-alpha inhibitor proteins (I α Ip) are a family of structurally related serine protease inhibitors found at relatively high concentrations (400-800 mg/L) in human plasma. I α Ip is a large, multi-component complex that functions as a trypsin-type protease inhibitor. Unlike other inhibitor molecules, this family of inhibitors consists of a combination of polypeptide chains (light and heavy chains) covalently linked uniquely by a chondroitin sulfate chain. The heavy chains of Inter-alpha proteins (H1, H2 and H3) are also called Hyaluronic acid (HA) binding proteins. The major forms found in human plasma are inter-alpha-inhibitor (I α I), which consists

of two heavy chains (H1 & H2) and a single light chain (L), and pre-alpha-inhibitor (PaI), which consists of one heavy (H3) and one light chain (L). The light chain (also termed bikunin (*Bikunitz inhibitor*, having two Kunitz domains) is known to broadly inhibit plasma serine proteases. The complex has been shown to be important in the inhibition of an array of proteases, including neutrophil elastase, plasmin, trypsin, chymotrypsin, cathepsin G, and acrosin.

IaI and PaI have also been found to be complexed with H4, another heavy chain of IaIp proteins. Certain embodiments of IaIp compositions according to the invention contain H4 in complex with PaI, IaI, or both PaI and IaI.

Without wishing to be bound by any scientific theories, we speculate that heavy chains of IaIp, after being released from the complex, bind HA preventing HA from binding its receptor, CD44. In the absence of heavy chains of IaIp, HA will bind to CD44 and trigger the secretion of pro-inflammatory factors, for example, TNF-alpha, and cause inflammation. Meanwhile, the light chains of IaIp, once released from the complex exhibit anti-protease activity.

"IaIp composition" refers to a preparation of IaIp proteins, including IaI and PaI in physiological proportions. Physiological proportions, as used herein is intended to include proportions found in a person or animal that is not suffering from an infection or condition, and/or the ratio of IaI to PaI that appears naturally in human plasma. Physiological proportions are usually from between about 60% to about 80% IaI and between about 40% to about 20% PaI. Physiological proportions may vary from these ranges due to normal variations in genetic makeup of subjects.

As used herein, "mixture of inter-alpha inhibitor protein (IaI) and pre-alpha protein (PaI)" refers to a composition containing both the IaI and PaI complexes. The mixture may also contain buffers, salts, or other components that are used to isolate the IaIp complex. In certain aspects, the IaI and the PaI are present in the mixture in a physiological proportion.

IaI and PaI present in the plasma fraction have an apparent molecular weight of between about 60,000 to about 280,000 kDa. Molecular weight may be determined by sodium dodecyl sulfate polyacrylamide gel electrophoresis.

“Half-life,” as used herein, refers to half of the amount of time that the administered IaIp is active upon administration. IaIp compositions according to the invention have a half-life of, for example, greater than about 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 7.5, or 10 hours. In a preferred embodiment, the IaIp composition has a half-life of greater than about 5 hours. In a particularly preferred embodiment, the IaIp composition has a half-life of greater than about 10 hours. Longer half-lives are preferred, for example, because fewer doses are required to be administered to a subject over time.

The IaIp compositions of the invention may have a high trypsin inhibitory specific activity. The trypsin inhibitory specific activity of the IaIp compositions according to the invention may range from between about 1000 to about 2000 IU/mg. Preferably the trypsin inhibitory specific activity is above 1200 IU/mg and even more preferably above 1500 IU/mg. The trypsin inhibitory specific activity may be measured, for example, by the trypsin inhibitory assay using L-BAPA as a substrate. See, HU Bergmeyer, ed: vol 5, 3rd ed. 119 (1984) Verlag Chemie, Weinheim: Chromogenic substrate for the assay of trypsin: R. Geiger, H. Fritz, Methods of Enzymatic Analysis.

A composition of IaIp may be a mixture of inter-alpha inhibitor protein (IaI) and pre-alpha protein (PaI), wherein the IaI and the PaI are present in said mixture in a physiological proportion comprising a light chain of inter-alpha inhibitor protein associated with at least one of three heavy chains H1, H2 and H3. A composition according to the invention may also have a light chain of inter-alpha inhibitor protein associated with at least one of four heavy chains H1, H2, H3 and H4. Examples of each protein in the IaIp complex are as follows: Bikunin GenBank accession number: AAB84031, P02760; H1 GenBank accession number: P19827, NP_002206; H2 GenBank accession number: NP_002207, P19823; H3 GenBank accession number: NP_002208; H4 GenBank accession number: Q14624, NP_002209, which are each incorporated herein by reference in their entirety.

As used herein “blood plasma-derived” refers to being originally isolated or purified from blood plasma. That is, the natural environment of the composition is blood plasma.

As used herein, “a plasma fraction” is a fraction from an isolation or purification step, for example, chromatography, that was originally derived from blood plasma. Plasma fractions

according to the invention may be for example, a side fraction obtained from the purification of clotting factor IX, a side fraction from the purification of a prothrombin complex concentrate, a cryosupernatant resulting from cryoprecipitation (described in Hoffer et al., Journal of Chromatography B 669 (1995) 187-196) of blood plasma, or cryo-poor plasma cryo-poor plasma
5 is used interchangeably with cryosupernatant herein. The cryo-poor plasma is the supernatant obtained from cryoprecipitation.

An example of a side fraction according to the invention is one obtained from the purification of clotting factor IX. A mixture of IaI/PaI has been shown to be present in side-fractions generated during the purification of factor IX (FIX). The method of obtaining the side
10 fraction obtained from the purification of clotting factor IX is described in Hoffer et al., Journal of Chromatography B 669 (1995) 187-196, which is hereby incorporated by reference in its entirety. Other examples of side fractions include a side fraction from FIX purification or a side fraction from the purification of a prothrombin complex concentrate, as is described in D. Josic et al., Thrombosis Research 100 (2000) 433-441, which is hereby incorporated by reference in its
15 entirety; and a side fraction isolated as a cryosupernatant resulting from cryoprecipitation of blood plasma. (For example, suitable cryoprecipitation methods are described in Hoffer et al., Journal of Chromatography B 669 (1995) 187-196.) Other plasma fractions that may be useful to purify IaIp complexes from blood include strong anion-exchange fractions and monolith chromatographic fractions, which are described below in the examples.

20 Plasma fractions, according to the invention, may be from human, primate, bovine, porcine, feline, or canine sources.

As used herein, the term "obtaining" includes purchasing, synthesizing, isolating or otherwise acquiring one or more of the substances used in practicing the invention.

Thus, "obtaining blood," as used herein, includes acquiring blood, for example, from
25 human, primate, bovine, porcine, feline, or canine sources. The blood may be acquired and/or purchased from, for example, blood banks, hospitals, hospices, private companies, research foundations, or any other source of blood.

“Obtaining blood plasma,” as used herein, includes acquiring blood plasma, for example, from human, primate, bovine, porcine, feline, or canine sources. The blood plasma may be acquired and/or purchased from, for example, blood banks, hospitals, hospices, private companies, research foundations, or any other source of blood. Alternately, blood plasma may also be isolated from blood once blood is obtained. Suitable methods of isolating blood plasma include gravity and centrifugation.

As used herein, “obtaining a side fraction obtained from the purification of clotting factor IX,” includes acquiring and/or purchasing side fractions obtained from the purification of clotting factor IX from for example, a company or hospital that routinely purifies factor IX.

As used herein, “obtaining a side fraction from the purification of a prothrombin complex concentrate,” includes acquiring and/or purchasing the side fractions, for example, from a company, research organization or hospital that purifies prothrombin complex.

As used herein, “obtaining a cryosupernatant resulting from cryoprecipitation of blood plasma,” includes acquiring and/or purchased a cryosupernatant, for example, from a hospital, research organization, or company that cryoprecipitates blood plasma in a manner suitable for use in the invention.

“Solid support” refers to a solid material which can be derivatized with, or otherwise attached to, a capture reagent. Exemplary solid supports include probes, microtiter plates and chromatographic resins.

“Adsorption” refers to detectable non-covalent binding of an analyte to an adsorbent or capture reagent. An adsorbent surface refers to a surface to which is bound an adsorbent (also called a “capture reagent” or an “affinity reagent”). An adsorbent is any material capable of binding an analyte (e.g., a target polypeptide or nucleic acid).

A chromatographic adsorbent refers to a material typically used in chromatography. Chromatographic adsorbents include, for example, ion exchange materials, metal chelators (e.g., nitriloacetic acid or iminodiacetic acid), immobilized metal chelates, hydrophobic interaction adsorbents, hydrophilic interaction adsorbents, dyes, simple biomolecules (e.g., nucleotides,

amino acids, simple sugars and fatty acids) and mixed mode adsorbents (e.g., hydrophobic attraction/electrostatic repulsion adsorbents).

A biospecific adsorbent refers to an adsorbent comprising a biomolecule, e.g., a nucleic acid molecule (e.g., an aptamer), a polypeptide, a polysaccharide, a lipid, a steroid or a conjugate of these (e.g., a glycoprotein, a lipoprotein, a glycolipid, a nucleic acid (e.g., DNA)-protein conjugate). In certain instances the biospecific adsorbent can be a macromolecular structure such as a multiprotein complex, a biological membrane or a virus. Examples of biospecific adsorbents are antibodies, receptor proteins and nucleic acids. Biospecific adsorbents typically have higher specificity for a target analyte than chromatographic adsorbents.

“Eluent” or “wash solution” refers to an agent, typically a solution, which is used to affect or modify adsorption of an analyte to an adsorbent surface and/or remove unbound materials from the surface. The elution characteristics of an eluant can depend, for example, on pH, ionic strength, hydrophobicity, degree of chaotropism, detergent strength and temperature.

“Analyte” refers to any component of a sample that is desired to be detected. The term can refer to a single component or a plurality of components in the sample.

The terms “polypeptide,” “peptide,” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residues are analogs or mimetics of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers. Polypeptides can be modified, e.g., by the addition of carbohydrate residues to form glycoproteins. The terms “polypeptide,” “peptide” and “protein” includes glycoproteins, as well as non-glycoproteins.

“Immunoassay” is an assay that uses an antibody to specifically bind an antigen (e.g., an IgA complex). The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the antigen.

“Antibody” refers to a polypeptide ligand substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof, which specifically binds and recognizes an epitope (e.g., an antigen). The recognized immunoglobulin genes include the kappa and lambda light chain constant region genes, the alpha, gamma, delta, epsilon and mu heavy chain constant

region genes, and the myriad immunoglobulin variable region genes. Antibodies exist, *e.g.*, as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. This includes, *e.g.*, Fab' and F(ab')₂ fragments. Antibodies includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

The phrase "specifically (or selectively) binds" to an antibody or "specifically (or selectively) immunoreactive with," when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies raised to the IαIp complex from a specific species such as rat, mouse, or human can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with IαIp complex and not with other proteins, except for polymorphic variants and alleles of the IαIp complex. This selection may be achieved by subtracting out antibodies that cross-react with the IαIp complex molecules from other species. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (*see, e.g.*, Harlow & Lane, *Antibodies, A Laboratory Manual* (1988), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity). Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

"Functionally equivalent" as used herein refers to any protein capable of exhibiting a substantially similar *in vivo* or *in vitro* activity as the IαIp proteins described herein, *e.g.*, effecting a decrease in sepsis.

IαIp product structure and purity may be confirmed by, for example, HPLC or other chromatographic method known to one of skill in the art.

As used herein, "I α Ip complex" is intended to encompass all naturally occurring biologically active variants of the I α Ip proteins, including proteins containing deletions, insertions, additions, and substitutions. A "natural variant" of an I α Ip protein is defined as a peptide obtained from plasma having a sequence that is altered by one or more amino acids. The
5 variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. In other embodiments, a variant may have "nonconservative" changes, e.g., replacement of a glycine with a tryptophan. Similar variations may also include amino acid deletions or insertions, or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted
10 without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTAR software.

A "deletion" is defined as a change in either amino acid or nucleotide sequence in which one or more amino acid or nucleotide residues, respectively, are absent.

An "insertion" or "addition" is that change in an amino acid or nucleotide sequence
15 which has resulted in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the naturally occurring I α Ip complex.

A "substitution" results from the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

The term "biologically active" refers to having structural, regulatory or biochemical
20 functions of a naturally occurring I α Ip complex. Likewise, "immunologically active" defines the capability of the natural, recombinant or synthetic I α Ip complex, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "derivative" as used herein refers to the chemical modification of a nucleic acid
25 encoding I α Ip complex or the encoded I α Ip complex. Illustrative of such modifications would be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would encode a polypeptide which retains essential biological characteristics of natural I α Ip complex.

“Purifying,” as used herein, refers to steps or processes of removing unwanted or contaminating proteins or components from IaIp to produce a purified IaIp complex. For example, a plasma fraction containing IaI and PaI in physiological proportion may be run through a series of chromatography steps to purify the IaIp composition.

5 “Isolating,” as used herein refers to producing a plasma fraction from blood plasma, which contains IaI and PaI in physiological proportions. For example, isolating a plasma fraction may be achieved in accordance with the invention by chromatographing blood plasma. Isolated refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered “by the hand of man” from its natural state. For
10 example, an isolated polypeptide or protein could be a component of blood plasma, or could be contained within a cell and be considered “isolated” because that blood plasma or particular cell may not be the original environment of the polypeptide.

Chromatographing, as used herein, may include anion-exchange chromatography. Anion-exchange chromatography may be particle-based, for example, DEAE Sepharose, DEAE
15 Sephadex A50, Toyopearl DEAE, TMAE Fractogel, DEAE Fractogel, or Q-Sepharose. Anion-exchange chromatography may also be by monolithic support, for example, CIM with immobilized anion-exchange ligands such as DEAE-CIM or Q-CIM. SEPHAROSE is a trade name of Pharmacia, Inc. of New Jersey for a high molecular weight substance for the separation by gel filtration of macromolecules. Anion exchange columns have two components, a matrix
20 and a ligand. The matrix can be, for example, cellulose, dextrans, agarose or polystyrene. The ligand can be diethylaminoethyl (DEAE), polyethyleneimine (PEI) or a quaternary ammonium functional group. The strength of an anion exchange column refers to the state of ionization of the ligand. Strong anionic exchange columns, such as those, having a quaternary ammonium ligand, bear a permanent positive charge over a wide pH range. In weak anion exchange
25 columns, such as DEAE and PEI, the existence of the positive charge depends on the pH of the column. Strong anion exchange columns such as Q Sepharose FF, or metal-chelating Sepharose (e. g., Cu²⁺-chelating Sepharose) are preferred. Anion exchange columns are generally loaded with a low-salt buffer at a pH above the pI of α -glucosidase.

The IaIp compositions of the invention are preferably from between about 85% to about 100% pure. As used herein, the term "pure" refers to the IaIp composition that is removed from its natural environment, isolated or separated, and is at least between about 85% to about 100% free, preferably 90% free, and more preferably 95% free from other components with which it is naturally associated. In preferred embodiments, a substantially purified protein will constitute more than 85%, 87.5%, 90%, 92.5%, 95%, 99% or even more of the proteins in the composition.

A peptide, polypeptide or protein that is "purified to homogeneity," as applied to the present invention, means that the peptide, polypeptide or protein has a level of purity where the peptide, polypeptide or protein is substantially free from other proteins and biological components. Any suitable materials and methods can be used to perform the isolation step or steps of blood plasma to obtain purified IaIp.

Typically, preparation of IaIp involves isolation of the sample and collection of fractions determined to contain the proteins of interest. Methods of isolation include, for example, solid phase extraction, chromatography, for example anion-exchange chromatography, size exclusion chromatography, ion exchange chromatography, heparin chromatography, affinity chromatography, sequential extraction, gel electrophoresis and liquid chromatography. Preparation may also include purifying, which may include the steps of chromatography, for example, ion exchange chromatography, heparin chromatography, affinity chromatography, sequential extraction, gel electrophoresis and liquid chromatography.

In one embodiment of the invention, a sample can be purified twice by anion exchange chromatography. Anion exchange chromatography allows purification of the proteins in a sample roughly according to their charge characteristics. For example, a Q anion-exchange resin can be used (*e.g.*, Q HyperD F, Biosepra), and a sample can be sequentially eluted with eluants having different pH's. Anion exchange chromatography allows separation of biomolecules in a sample that are more negatively charged from other types of biomolecules. Proteins that are eluted with an eluant having a high pH is likely to be weakly negatively charged, and a fraction that is eluted with an eluant having a low pH is likely to be strongly negatively charged. Thus, in addition to reducing complexity of a sample, anion exchange chromatography separates proteins according to their binding characteristics.

In yet another embodiment, a sample can be further purified by heparin chromatography. Heparin chromatography allows further purification of the IaIp complexes in a sample also on the basis of affinity interaction with heparin and charge characteristics. Heparin, a sulfated mucopolysaccharide, will bind IaIp complexes with positively charged moieties and a sample
5 can be sequentially eluted with eluants having different pH's or salt concentrations. IaIp complexes eluted with an eluant having a low pH are more likely to be weakly positively charged. IaIp complexes eluted with an eluant having a high pH are more likely to be strongly positively charged. Thus, heparin chromatography also reduces the complexity of a sample and separates IaIp complexes according to their binding characteristics.

10 IaIp complexes may be may be captured with capture reagents immobilized to a support, such as any biochip, a multiwell microtiter plate, a resin, or nitrocellulose membranes that are subsequently probed for the presence of proteins. In particular, the IaIp complexes of this invention may be captured on Surface-Enhanced Laser Desorption/Ionization (SELDI) protein
15 biochips. Capture can be on a chromatographic surface or a biospecific surface. Any of the SELDI protein biochips comprising reactive surfaces can be used to capture and detect the IaIp complexes of this invention. These biochips can be derivatized with the antibodies that specifically capture the IaIp complexes, or they can be derivatized with capture reagents, such as protein A or protein G that bind immunoglobulins. Then the IaIp complexes can be captured in
20 solution using specific antibodies and the captured IaIp complexes isolated on chip through the capture reagent.

Various methods for quantifying the degree of purification of proteins, polypeptides, or peptides will be known to those of skill in the art in light of the present disclosure. These include, for example, determining the specific protein activity of a fraction, or assessing the number of polypeptides within a fraction by gel electrophoresis.

25 In addition to those techniques described in detail herein below, various other techniques suitable for use in protein purification will be well known to those of skill in the art. These include, for example, precipitation with ammonium sulfate, PEG, antibodies and the like or by heat denaturation, followed by centrifugation; chromatography steps such as ion exchange, gel filtration, reverse phase, hydroxylapatite, lectin affinity, immunoaffinity chromatography and

other affinity chromatography steps; isoelectric focusing; gel electrophoresis, HPLC; and combinations of such and other techniques. Furthermore, if viewed as desirable, additional purification steps can be employed using approaches that are standard in this art. These approaches are fully able to deliver a highly pure preparation of the protein.

5 In other embodiments, gel chromatography, or molecular sieve chromatography may be used. Gel chromatography is a special type of partition chromatography that is based on molecular size. The theory behind gel chromatography is that the column, which is prepared with tiny particles of an inert substance that contain small pores, separates larger molecules from smaller molecules as they pass through or around the pores, depending on their size. As long as
10 the material of which the particles are made does not adsorb the molecules, the sole factor determining rate of flow is the size. Hence, molecules are eluted from the column in decreasing size, so long as the shape is relatively constant. Gel chromatography is unsurpassed for separating molecules of different size because separation is independent of all other factors such as pH, ionic strength, temperature, etc. There also is virtually no adsorption, less zone spreading
15 and the elution volume is related in a simple matter to molecular weight.

 In other embodiments, affinity chromatography may be used. Affinity chromatography is a chromatographic procedure that relies on the specific affinity between a substance to be isolated and a molecule that it can specifically bind to. This is, for example, a receptor-ligand type interaction. The column material may be synthesized by covalently coupling one of the
20 binding partners to an insoluble matrix. The column material may then be able to specifically adsorb the substance from the solution. Elution occurs, for example, by changing the conditions to those in which binding will not occur (e.g., alter pH, ionic strength, and temperature.).

 The methods described herein are amenable for large-scale production, and result in proteins including, the IaIp complex and H4 in a form suitable for therapeutic administration.

25 Any of the isolation or purification steps of the processes described herein may be repeated to obtain a higher degree of purity, if desired. The chromatography steps may be in batch or in column format.

Preferred processes for producing a blood plasma-derived IaIp composition of the invention include isolating from blood plasma a plasma fraction containing IaI and PaI, wherein the IaI and PaI are present in a physiological proportion; and purifying the plasma fraction to obtain an IaIp composition with a purity of from between about 85% to about 100% pure.

5 Processes according to the invention may also include further purification of the plasma fraction, for example, by passing to a heparin affinity column and collecting the flow through (unbound) fraction.

Processes useful in practicing the invention may also include virus inactivating the plasma fraction and/or the purified IaIp before and/or after the purification and isolation steps.
10 Typical processes include virus inactivating by a solvent/detergent treatment or thermal inactivation. Thermal inactivations, or pasteurization of the compositions of the invention may be for example at a temperature of between about 55° to about 65°C or dry heat at 70 to 120°C.

For solvent detergent treatments, a specific combination of solvent and detergents, such as, 0.3% tri-n-butylphosphate (TnBP) combined with 1% Tween-80, at 24°C for 6 hours, is
15 effective to inactivate enveloped viruses (Horowitz et al (1985) Transfusion 25, pp. 516-522). Alternatively, pasteurization of the fraction or purified IaIp at 55-65 °C in the presence of stabilizers may be sufficient to inactivate any viruses present.

During the purification or isolation step(s) stabilizers may be added. The final composition of IaIp may also contain stabilizers. For example, suitable stabilizers
20 include albumin, polyethylene glycol, alpha, alpha-trehalose, amino acids, salts, glycerol, omega-amino acids such as lysine, polylysine, arginine, epsilon amino caproic acid and tranexamic acid, sugars, such as sucrose, or combinations thereof.

The IaIp proteins and compositions of the invention may be used to treat a human disease. Such diseases include, for example, acute inflammatory diseases, sepsis, severe shock,
25 septic shock, rheumatoid arthritis, cancer, cancer metastasis, preterm labor and infectious diseases. The IaIp compositions used to treat disease may be made by isolating from blood plasma a plasma fraction containing IaI and PaI, wherein the IaI and PaI are present in a

physiological proportion; and purifying the plasma fraction to obtain an I α Ip composition with a purity of I α Ip ranging from about 85% to about 100% pure.

The invention also encompasses pharmaceutical compositions of I α Ip. Pharmaceutical compositions of I α Ip may be any of the I α Ip compositions described herein in a therapeutically effective amount with a pharmaceutically acceptable carrier.

For example, a pharmaceutical composition according to the invention may be a therapeutically effective amount of an I α Ip composition which is a mixture of inter-alpha inhibitor protein (I α I) and pre-alpha protein (PaI), wherein the I α I and the PaI are present in said mixture in a physiological proportion ranging from about 85% to about 100% pure. A pharmaceutical composition may also be a therapeutically effective amount of an I α Ip composition which is a mixture of inter-alpha inhibitor protein (I α I) and pre-alpha protein (PaI), wherein the I α I and the PaI are present in said mixture in a physiological proportion comprising a light chain of inter-alpha inhibitor protein associated with at least one of three heavy chains H1, H2 and H3 or H1, H2, H3 and H4. A pharmaceutical composition may also be a therapeutically effective amount of an I α Ip composition which is a mixture of inter-alpha inhibitor protein (I α I) and pre-alpha protein (PaI), wherein the I α I and the PaI are present in said mixture in a physiological proportion and having a high trypsin inhibitory specific activity. A pharmaceutical composition may also be a therapeutically effective amount of an I α Ip composition which is a mixture of inter-alpha inhibitor protein (I α I) and pre-alpha protein (PaI), wherein the I α I and the PaI are present in said mixture in a physiological proportion and having a half life of greater than one hour, five hours, or ten hours.

The term "pharmaceutically acceptable carrier or adjuvant" refers to a carrier or adjuvant that may be administered to a subject, together with a composition of this invention, and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver a therapeutic amount of the composition.

Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as d- α -tocopherol polyethyleneglycol 1000 succinate, surfactants used in pharmaceutical dosage

forms such as Tweens or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, 5 sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Cyclodextrins such as α -, β -, and γ -cyclodextrin, or chemically modified derivatives such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl- β -cyclodextrins, or other 10 solubilized derivatives may also be advantageously used to enhance delivery of compositions described herein.

The pharmaceutical compositions of this invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir, preferably by oral administration or administration by injection.

15 The pharmaceutical compositions of this invention may contain any conventional non-toxic pharmaceutically-acceptable carriers, adjuvants or vehicles. In some cases, the pH of the formulation may be adjusted with pharmaceutically acceptable acids, bases or buffers to enhance the stability of the formulated composition or its delivery form. The term parenteral as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, 20 intraarterial, intrasynovial, intrasternal, intrathecal, intralesional and intracranial injection or infusion techniques. Suitable methods of administration may be as a tablet, capsule, or by intravenous injection. Injectable forms of administration are particularly preferred.

The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example, as a sterile injectable aqueous or oleaginous suspension. This suspension may be 25 formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride

solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, or carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms such as emulsions and or suspensions. Other commonly used surfactants such as Tweens or Spans and/or other similar emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, emulsions and aqueous suspensions, dispersions and solutions. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions and/or emulsions are administered orally, the active ingredient may be suspended or dissolved in an oily phase is combined with emulsifying and/or suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

The pharmaceutical compositions of this invention may also be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing a composition of this invention with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the active components. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.

Topical administration of the pharmaceutical compositions of this invention is useful when the desired treatment involves areas or organs readily accessible by topical application. For application topically to the skin, the pharmaceutical composition should be formulated with a

suitable ointment containing the active components suspended or dissolved in a carrier. Carriers for topical administration of the compositions of this invention include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene composition, emulsifying wax and water. Alternatively, the pharmaceutical composition can be formulated with a suitable lotion or cream containing the active composition suspended or dissolved in a carrier with suitable emulsifying agents. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water. The pharmaceutical compositions of this invention may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation. Topically-transdermal patches are also included in this invention.

The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

Methods of treating an inflammation related disorders, for example, rheumatoid arthritis, septic and septic shock, head trauma/injury and meningitis, inflammatory bowel diseases (Crohn's Disease), chronic obstructive pulmonary disease, rhinitis; cancer, preterm labor, or an infectious disease in a subject according to the invention may include administering a therapeutically effective amount of an IaIp composition that is produced according to the methods of the invention.

Compositions herein are administered in a dosage ranging from about 1 to 50 mg/kg of body weight, preferably dosages between 500 mg and 1000 mg/dose, every 4 to 120 hours, or according to the requirements of the particular drug. The methods herein contemplate administration of an effective amount of composition to achieve the desired or stated effect. Typically, the pharmaceutical compositions of this invention will be administered from about 1 to about 6 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with

the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. A typical preparation will contain from about 5% to about 95% active composition (w/w). Alternatively, such preparations contain from about 20% to about 80% active composition.

5 Lower or higher doses than those recited above may be advantageous. Specific dosage and treatment regimens for any particular subject will depend upon a variety of factors, including the activity of the specific composition employed, the age, body weight, general health status, sex, diet, time of administration, rate of excretion, drug combination, the severity and course of the disease, condition or symptoms, the subject's disposition to the disease, condition or
10 symptoms, and the judgment of the treating physician.

Upon improvement of a condition, a maintenance dose of a composition or combination of this invention may be administered, if necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained when the symptoms have been alleviated to the desired level,
15 treatment should cease. Subjects may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms. Improvement of the condition may also be judged based upon the level of IaIp in the liver. If normal physiological levels of IaIp are found in the liver and the patient's symptoms as judged by the subject or the treating physician are judged to be improved, the subject may be treated with a maintenance dose.

20 When the compositions of this invention comprise a combination of an IaIp composition and one or more additional therapeutic or prophylactic agents, both the composition and the additional agent should be present at dosage levels of between about 1 to 100%, and more preferably between about 5 to 95% of the dosage normally administered in a monotherapy regimen. The additional agents may be administered separately, as part of a multiple dose
25 regimen, from the compositions of this invention. Alternatively, those agents may be part of a single dosage form, mixed together with the compositions of this invention in a single composition.

Methods for treating a subject for acute inflammatory disease, sepsis, severe shock, septic shock, rheumatoid arthritis, cancer, cancer metastasis, infectious disease, and/or preterm labor,

include the steps of determining the pre-treatment level of one or more of IaI, PaI, IaIp, H3, H4, H1, H2, and LC; and administering a therapeutically effective amount of IaIp to the subject. Pre-treatment levels of IaI, PaI, IaIp, H3, H4, H1, H2, and LC are the levels of the proteins in the subject prior to the first administration of IaIp or any of the IaIp complex proteins. Post-treatment levels are the levels of IaIp measured after administration of IaIp. The methods of the invention include determining the post-treatment levels of one or more of IaI, PaI, IaIp, H3, H4, H1, H2, and LC after an initial period of treatment with IaIp. A modulation in the level of IaIp is an indication that the treatment is producing a favorable clinical response. The initial period of treatment may be the time required to achieve a steady-state plasma concentration of the IaIp.

The level of IaI, PaI, IaIp, H3, H4, H1, H2, and/or LC may be determined, for example, by immunological methods. For example, IaIp complexes can be detected and/or measured by a variety of detection methods including for example, gas phase ion spectrometry methods, optical methods, electrochemical methods, atomic force microscopy, radio frequency methods, surface plasmon resonance, ellipsometry immunological, and atomic force microscopy methods.

In another embodiment, an immunoassay can be used to detect and analyze IaIp complexes in a sample. This method comprises: (a) providing an antibody that specifically binds to an IaIp complex; (b) contacting a sample with the antibody; and (c) detecting the presence of a complex of the antibody bound to the IaIp complex in the sample. Suitable antibodies for use in the methods of the invention include, MA b 69.31, MA b 69.26, anti-IaIp polyclonal antibody (R16 or R20), and anti-bikunin monoclonal or polyclonal antibody.

An immunoassay is an assay that uses an antibody to specifically bind an antigen (*e.g.*, an IaIp complex). The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the antigen. The phrase "specifically (or selectively) binds" to an antibody or "specifically (or selectively) immunoreactive with," when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an

antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies raised to an IaIp complex from specific species such as rat, mouse, or human can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with that IaIp complex and not with other proteins, except for polymorphic variants and alleles of the IaIp complex. This selection may be achieved by subtracting out antibodies that cross-react with the IaIp complex molecules from other species.

Subjects suitable for treatment with IaIp may be identified as having inflammation, trauma/injury, tumor invasion, tumor metastasis, sepsis, septic shock, or an infectious disease. The subject may be self-identified or diagnosed by a medical practitioner as having inflammation, tumor invasion, tumor metastasis, sepsis, septic shock, or an infectious disease. The subjects may be primates, humans, or other animals.

Methods also include the co-administration of other therapeutic agents. For example, the additional therapeutic agents may be anticancer agents, anti-inflammatory agents, anti-coagulants or immunomodulators. For example, dideoxynucleosides, e.g. zidovudine (AZT), 2',3'-dideoxyinosine (ddI) and 2',3'-dideoxycytidine (ddC), lamivudine (3TC), stavudine (d4T), and TRIZIVIR (abacavir + zidovudine + lamivudine), nonnucleosides, e.g., efavirenz (DMP-266, DuPont Pharmaceuticals/Bristol Myers Squibb), nevirapine (Boehringer Ingelheim), and delaviridine (Pharmacia-Upjohn), TAT antagonists such as Ro 3-3335 and Ro 24-7429, protease inhibitors, e.g., indinavir (Merck), ritonavir (Abbott), saquinavir (Hoffmann LaRoche), nelfinavir (Agouron Pharmaceuticals), 141 W94 (Glaxo-Wellcome), atazanavir (Bristol Myers Squibb), amprenavir (GlaxoSmithKline), fosamprenavir (GlaxoSmithKline), tipranavir (Boehringer Ingelheim), KALETRA (lopinavir + ritonavir, Abbott), and other agents such as 9-(2-hydroxyethoxymethyl)guanine (acyclovir), interferon, e.g., alpha-interferon, interleukin II, and phosphonoformate (Foscarnet), or entry inhibitors, e.g., T20 (enfuvirtide, Roche/Trimeris) or UK-427,857 (Pfizer), levamisole or thymosin, cisplatin, carboplatin, docetaxel, paclitaxel, flurouracil, capecitabine, gemcitabine, irinotecan, topotecan, etoposide, mitomycin, gefitinib, vincristine, vinblastine, doxorubicin, cyclophosphamide, celecoxib, rofecoxib, valdecoxib, ibuprofen, naproxen, ketoprofen, dexamethasone, prednisone, prednisolone, hydrocortisone, acetaminophen, misonidazole, amifostine, tamsulosin, phenazopyridine, ondansetron, granisetron, alosetron, palonosetron, promethazine, prochlorperazine, trimethobenzamide,

aprepitant, diphenoxylate with atropine, and/or loperamide. Anti-coagulants such as Anti-thrombin III, activated Protein C and protease inhibitors such as furin inhibitors

Methods also include predicting a response to an IaIp therapy by assaying a sample obtained from a subject to detect the level of one or more of IaI, PaI, IaIp, H3, H4, H1, H2, and
5 LC; wherein the detected levels identifies a subject that may respond favorably to IaIp therapy. For example, a decrease in a detectable level of IaI and/or PaI indicates that a subject may benefit from the administration of IaIp.

Methods also include monitoring the progress of a subject being treated with an IaIp therapy by determining the pre-treatment level of one or more of IaI, PaI, IaIp, H3, H4, H1, H2,
10 and LC; administering a therapeutically effective amount of IaIp to the subject; and determining the level of one or more of the levels in the subject after an initial period of treatment with the IaIp, wherein an increase of the level in the subject following treatment with IaIp indicates that the subject is likely to have a favorable clinical response to treatment with IaIp.

Kits for IaIp therapy may include one ore more of IaI, PaI, IaIp, H3, H4, H1, H2, and
15 LC; and instructions for therapeutic use. Kits are also contemplated that have the IaIp compositions described herein and instructions for use. For example, a kit may have IaI, PaI, and instructions that provide information regarding dosage, form of administration, and storage conditions.

A container including IaIp and a label or package insert with instructions for
20 administering the IaIp to a subject are also contemplated. The instructions may provide instructions for dosage, form of administration, and storage conditions.

EXAMPLES

It should be appreciated that the invention should not be construed to be limited to the
25 examples which are now described; rather, the invention should be construed to include any and all applications provided herein and all equivalent variations within the skill of the ordinary artisan.

EXAMPLE 1**Characterization of H4 As A Part of the I α I β Complex**

SELDI-TOF mass spec analysis indicated that heavy chain 4 (H4) is also present in both 125 kDa band (P α I) and 250 kDa band (I α I) (data not shown). The presence of H4 in the 250 kDa band is more pronounced. This suggests that another complex protein other than I α I (H1+H2+LC) and P α I (H3+LC) might be present in some compositions of the invention. So far, H4 has not been described in the complexed form. Free H4 is smaller than 125 or 250 kDa. Due to the mass spec results, we think that H4 might be in complex with bikunin (the light chain) or something else.

EXAMPLE 2**Animal model of sepsis**

Male Sprague-Dawley rats (275-325g) were housed in a temperature-controlled room on a 12-h light/dark cycle and fed a standard Purina rat chow diet. Prior to the induction of sepsis, rats were fasted overnight but allowed water *ad libitum*. Rats were anesthetized with isoflurane inhalation and the ventral neck, abdomen and groin were shaved and washed with 10% povidone iodine. A 2-cm midline abdominal incision was performed. The cecum was exposed, ligated just distal to the ileocecal valve to avoid intestinal obstruction, punctured twice with an 18-gauge needle, squeezed slightly to allow a small amount of fecal matter to flow from the holes, and returned to the abdominal cavity; the abdominal incision was then closed in layers. Sham-operated animals (i.e., control animals) underwent the same procedure with the exception that the cecum was neither ligated nor punctured. The animals were resuscitated with 3 ml/100g BW normal saline subcutaneously immediately after surgery. Animals were then anesthetized at various intervals after cecal ligation and puncture (CLP) or sham operation for collection of tissue samples. All experiments were performed in accordance with the National Institutes of Health guidelines for the use of experimental animals. This project was approved by the Institutional Animal Care and Use Committee of the North Shore-Long Island Jewish Research Institute.

Preparation and administration of human IaIp

Human IaIp (both IaI and PaI) was isolated as a by-product of a procedure designed for purifying coagulation factor VIII from human plasma. The procedure involves ion exchange and size-exclusion chromatography of cryoprecipitates. A purity of approximately 70% was achieved after the chromatographic separation. This IaIp containing preparation has few side effects in terms of toxicity, thrombogenicity, or hypotension. At 1, 5, 10 or 10 and 20 h after the onset of sepsis, the left femoral vein was cannulated with a polyethylene-50 tubing under isoflurane anesthesia. Human IaIp concentrates at a dose of 30 mg/kg BW or an equivalent volume of vehicle (normal saline, 1.5 ml/rat) were administered intravenously via the femoral catheter over 30 min at a constant infusion rate. Administration of human IaIp, in previous experiments, did not alter mean arterial pressure or heart rate during the period of infusion or thereafter (data not shown).

RNA extraction and determination of hepatic bikunin genes

The liver may be the major source of bikunin. Therefore, we measured bikunin mRNA expression in the liver. Total RNA was extracted from the liver by Tri-Reagent (Molecular Research Center, Cincinnati, OH). One hundred mg of liver tissue was homogenized in 1.5 ml Tri-Reagent, separated into aqueous and organic phases by chloroform addition, and centrifuged. RNA was precipitated from the aqueous phase by addition of isopropanol, and washed with ethanol. The pellet was dissolved in 0.1% DEPC-treated, deionized distilled water. RNA concentration and purity were determined by measuring the absorbance at 260 and 280 nm. Five µg of RNA from each tissue was reverse-transcribed in a 20 µl reaction volume containing 50 mM KCl, 10 mM Tris-HCl, 5 mM MgCl₂, 1 mM dNTP, 20 U RNase inhibitor, 2.5 mM oligo d(T)₁₆ primer and 50 U reverse transcriptase. The reverse transcription reaction solution was incubated at 42°C for 1 h, followed by heating at 95°C for 5 min. One µl cDNA was amplified with 0.15 µM each of 3' and 5' primers, specific for rat bikunin (633bp) (5' TGA CGA ATA TGC CAT TTT CC 3', 5' CCA CAG TAC TCC TTG CAC TCC 3') (accession No. S87544), rat glyceraldehydes-3-phosphate-dehydrogenase 7 (G3PDH) (24) (983bp) (5' TGA AGG TCG GTG TCA ACG GAT TTG GC 3', 5' CAT GTA GGC CAT GAG GTC CAC CAC 3') in 25 µl of PCR mixture containing 50 mM KCl, 10 mM Tris-HCl, 2 mM MgCl₂, 0.2 mM dNTP and 0.7 U

AmpliTaq DNA polymerase. PCR was carried out in a Bio-Rad thermal cycler. Following RT-PCR, 5 µl of the reaction mixture was electrophoresed in 1.2% TBE-agarose gel containing 0.22 µg/ml ethidium bromide. The gel was then developed and band intensities were normalized by G3PDH using the Bio-Rad image system (Hercules, CA).

5 Radioiodination of protein and determination of I αIp half-life

Purified IαIp was radioiodinated with Na¹²⁵I (Amersham, Arlington Heights, IL) using 1,3,4,6-tetrachloro- 3a-6a-diphenyl glycoluril (IODO-GEN iodination reagent; Pierce, Rockford, IL). The unincorporated ¹²⁵I was removed by applying the reaction mixture to an Excellulose GF-5 desalting column (Pierce). Radioactivity was determined in a gamma counter (Pharmacia-LKB, Piscataway, NJ). At 12 h after CLP or sham operation, the animals were anesthetized with isoflurane inhalation. A steady state of sedation was maintained with a subsequent intravenous injection of sodium pentobarbital (~30 mg/kg BW). Polyethylene-50 catheters were placed in the right jugular vein and left femoral artery, and a bolus injection of ¹²⁵I-labeled IαIp (~500,000 cpm/rat) was administered through the jugular cannula. The remaining radioactivity in the syringe was measured with a gamma counter, and the radioactivity counts were subtracted from the initial preinjection counts to determine the net injected radioactivity. Blood samples were collected immediately after the injection and then every 2 h for a period of 8 h for determining the half-life (t_{1/2}) of ¹²⁵I-IαIp in the circulation. The radioactivity (cpm) in each sample was measured with a gamma counter. The t_{1/2} was calculated according to Wu R, Zhou M, Cui X, et al: Ghrelin clearance is reduced at the late stage of polymicrobial sepsis. *Int J Mol Med*. 2003;12:777-782.

Survival study

CLP was performed as described above. At 1, 5, 10 or 10 and 20 h after CLP, human IαIp concentrate (30 mg/kg BW) or vehicle (normal saline, 1.5 ml/rat, at 1 h after CLP or 10 and 20 h after CLP) was infused intravenously. At 20 h after CLP, the necrotic cecum was excised and the abdominal cavity was washed twice by using 40 ml of warm, sterilized normal saline solution. The abdominal incision then was closed in layers. The procedure of cecal excision in CLP animals was performed to mimic the clinical situation in which septic focus should be removed

whenever possible. The animals then were allowed food and water *ad libitum* and were monitored for 10 days to record survival.

Statistical analysis

Results are expressed as means \pm SE. One-way analysis of variance (ANOVA) and Tukey's test were used to compare different groups of experimental animals. The survival rate was estimated by Kaplan-Meier method and compared by the log-rank test. Differences in values were considered significant if $P < 0.05$. 9

Alterations in bikunin mRNA expression after CLP

Bikunin is an active part of α_2 Ip. The liver is the major source of bikunin. Therefore, we chose bikunin mRNA expression in the liver to reflect the production of α_2 Ip. As shown in Figure 3, mRNA expression of bikunin in the liver did not change at 5 h after CLP, however, a 32% decrease was found at 20 h after CLP as compared with sham operated animals ($P < 0.05$).

Alterations in $t_{1/2}$ of 125 I- α_2 Ip after CLP

Half-life of α_2 Ip was estimated by measuring the changes of blood levels of radioactive labeled α_2 Ip injected at 12 h after the onset of sepsis. As indicated in Figure 4, the $t_{1/2}$ of 125 I- α_2 Ip was significantly increased from 5.6 ± 0.3 h to 11.8 ± 2.7 h ($P < 0.05$) after CLP.

Effects of α_2 Ip on survival rate

The survival rate after CLP and cecal excision with single time vehicle administration (at 1 h after CLP) was 75% at day 2 and decreased to 50% at days 5–10 (Figure 5). Administration of human α_2 Ip at 1 h after CLP, however, improved the survival rate to 92% throughout the 10-day observation period ($P < 0.05$; Figure 5). Although administration of human α_2 Ip at 5 or 10 h after CLP improved the survival rate to 64% and 73% respectively, these improvements were not statistically significant (Figure 6). The survival rate after CLP and cecal excision with two times vehicle administration (at 10 and 20 h after CLP) was 56% at day 2 and decreased to 44% at days 5–10 (Figure 7), which was not significantly different as compared with one time vehicle administration (Figure 5). Administration of human α_2 Ip at 10 and 20 h after CLP, however,

improved the survival rate to 81% throughout the 10-day observation period, which was significantly different as compared with vehicle group ($P < 0.05$; Figure 7).

Sepsis is a clinical syndrome characterized by systemic inflammation, coagulopathy, respiratory failure, myocardial dysfunction, renal insufficiency, and neurocognitive defects. It is generally assumed that this syndrome results from an excessive triggering of endogenous inflammatory mediators by the invading microorganisms. These mediators include substances released by activated monocytes, macrophages, endothelial cells and neutrophils such as cytokines, reactive oxygen species and proteases. In severe inflammatory response, various blood and tissue cells, including polymorphonuclear granulocytes, release lysosomal proteinases extracellularly and into the circulation. Such proteases as well as normally intracellular oxidizing agents produced during phagocytosis, can trigger tissue and organ damage and enhance the nonspecific proteolysis of plasma clotting and complement factors. The release of neutrophil proteinases, especially human leukocyte elastase, has been implicated in the progress of complications in subjects with sepsis. Their plasma levels are in close correlation with the severity of infection-induced inflammation and highly predictive of forthcoming organ failure.

During septic shock in humans, in addition to elevated activities of proteases, decreased plasma levels of I α Ip have been reported. Subjects with severely decreased concentrations of I α Ip have a higher mortality rate. Our results indicate that the gene expression of bikunin in the liver is significantly lower in the CLP animals than in the sham-operated animals. The expression of mRNA related to proteins in the inter-alpha-inhibitor family has been examined in various tissues in primates, pigs, and rodents. These studies indicate the genes which are the source of all members of the I α Ip family are primarily transcribed in the liver. Our results also show the gene expression of bikunin in other organs (i.e., intestine and kidneys) was not significantly altered at 5 or 20 h after CLP as compared to the shams (data not shown).

The significant downregulation observed only in the liver at 20 h after CLP suggests that this organ might be an important source of I α Ip and furthermore, in late stages of sepsis, bikunin gene expression in the liver is significantly decreased. Therapeutically, bikunin has been reported to have beneficial effects in humans as a prophylactic treatment to prevent pancreatitis after gastrectomy or to attenuate organ injury after cardiac surgery. Studies that examined the

effects of bikunin in an acute canine model of lethal *E. coli* bacteremia showed similar results with an improvement in hemodynamic variables and normalization of cardiac output and mean arterial pressure. Because the plasma half-life of bikunin is very short (approximately 10 min), it appears important to prolong the half-life of bikunin to maintain the sustained beneficial effects of this agent. Our results show that the half-life of IaIp in sham-operated animals is 5.6 h. When we injected radioactive labeled IaIp at 12 h after the onset of sepsis, we found the half-life of IaIp was prolonged to 11.8 h. These results indicate that IaIp clearance was significantly decreased during sepsis. However, even with decreased clearance the plasma levels of IaIp remain significantly lower in septic subjects. Our previous study has shown that administration of low purity IaIp early after the onset of sepsis (i.e., 1h post-CLP) maintained cardiac output and systemic oxygen delivery and increased systemic oxygen consumption and systemic oxygen extraction ratio. Moreover, IaIp downregulated TNF- α production and attenuated hepatocellular injury and lactic acidosis at 20 h after CLP. In addition, IaIp administration at 1 h after the onset of sepsis improved survival in septic animals.

Human IaIp proteins were isolated as a by-product of industrial scale plasma fractionation. The isolation method highly, simultaneously enriches the major plasma form of bikunin-containing proteins (IaI and PaI). Thus, in this preparation, a physiologic composition of plasma IaIp is obtained. The results of the mortality study performed indicate that administration of IaIp at 1 h after CLP improved the survival rate from 50% to 92% at 10 days after CLP and cecal excision. Although administration of human IaIp at 5 or 10 h after CLP improved the survival rate to 64% and 73% respectively, these improvements were not statistically significant.

Administration of human IaIp at 10 and 20 h after CLP, however, significantly improved the survival rate from 44% to 81%. Thus, IaIp appear to be a useful adjunct for improving survival during the progression of polymicrobial sepsis. In summary, bikunin gene expression in the liver decreased during sepsis and the half-life of IaIp increased from 5.6 ± 0.3 h to 11.8 ± 2.7 h, suggesting downregulation of bikunin in sepsis despite a decrease clearance. Administration of IaIp at 1 h after CLP improved the survival rate from 50% to 92%, whereas there was no significant improvement when IaIp was administrated at 5 or 10 h after CLP. However, double

injection of IaIp at 10 and 20 h after CLP increased the survival rate from 44% to 81%. Delayed but repeated administration of human IaIp improve survival after CLP.

EXAMPLE 3

5 Purification of IaIp

After application of dialyzed or ultra/diafiltrated eluate after solid-phase extraction with DEAE Sephadex A50, weakly bound components are eluted from DEAE-Sepharose FF column with 0.005 M sodium citrate/0.0055 M sodium phosphate buffer, pH 6.0 containing 0.28 M sodium chloride (described in Hoffer et al., Journal of Chromatography B 669 (1995) 187-196).

10 In the previous step, the column was washed with 0.005 M sodium citrate/0.0055 M sodium phosphate buffer, pH 6.0, containing 0.20 M sodium chloride.

After dialysis or ultrafiltration/diafiltration (UF/DF) against 0.005 M sodium phosphate buffer pH 7.0, the eluate was applied to hydroxylapatite column. The IaIp proteins do not bind to the column and are collected as flow through fraction. The contaminating proteins, mainly FII, FVII
15 and FX, can be eluted using a gradient with an increasing concentration of sodium phosphate buffer. The IaI/PaI fraction contains more than 90% of target proteins, mainly IaI and PaI.

EXAMPLE 4

Purification of IaIp from a Factor IX flow through fraction

20 Unbound proteins from Heparin Sepharose affinity chromatography (L. Hoffer et al., J. of Chromatography B), are applied to a DEAE-Sepharose FF anion-exchange column, After washing the column with a minimum of three column volumes of 0.005 M phosphate buffer, pH 7.0, IaIp/PaI containing fractions were eluted with 0.005 M phosphate buffer, pH 7.0, containing 0.55 M sodium chloride (elution buffer). The eluate contains about 30-40% IaI/PaI. After
25 dialysis or ultrafiltration/diafiltration (UF/DF) against 0.005 M sodium phosphate buffer, pH 7.0, the eluate from DEAE Sepharose FF was applied to a hydroxylapatite column. The IaI/PaI do

not bind to the column and are collected as a flow-through fraction. The contaminating proteins, mainly FII and FX can be eluted using a gradient with increasing concentration of sodium phosphate buffer. The purified IaI/PaI fraction contains more than 90% of the target proteins

5 EXAMPLE 5

Eluant from a solid-phase extraction of cryopoor plasma on DEAE-Sephadex A50 (L. Hoffer et al., J. of Chromatography B) or Q-Sephadex A50 (D. Josic et al., Thrombosis Research, cf. above) was applied to a DEAC-CIM tube monolith with a column volume of 80 mL (cf. K. Branovic et al., J of Chromatography A, 903 (2000) 21-32). The unbound fraction (flow-through) was collected. The column was subsequently washed with three column volume of 0.02 M Tris-HCl, pH 7.4 (equilibration buffer). Bound proteins were eluted in first step with 0.02 M Tris-HCl, pH 7.4, containing 0.35 Mol/L sodium chloride (Eluate 1) and in a second step with 0.02 M Tris-HCl, pH 7.4, containing 0.55 M sodium chloride (Elution 2). IaI/PaI are found in flow-through fractions and Eluate 1. Flow-through fractions contain about 35-45% IaI/PaI. The amount of these target proteins in Eluate 1 is between 20-30%. Fractions containing IaI/PaI were subjected to dialyses or ultrafiltration/diafiltration (UF/DF) against 0.005 M sodium phosphate buffer, pH 7 and applied to a hydroxylapatite column. IaIp do not bind to the column and are collected as a flow-through fraction. The flow-through fraction contains more than 90% of IaI/PaI.

Figures 8a and 8b describe exemplary purification schemes of IaIp in a flow diagram as represented by Examples 3-5.

EXAMPLE 6

Beneficial effects of highly purified IaIp in the animal studies.

Polymicrobial sepsis was induced by cecal ligation and pucture (CLP) in male Sprague-Dawley rats. Animals were fasted overnight before CLP was performed but were allowed water ad libitum. At the time of the experiment, the rats were anesthetized by methylflurane

inhalation, and a 2-cm ventral midline incision was performed. The caecum was exposed, ligated just distally to the ileocecal valve to avoid intestinal obstruction, punctured twice with an 18-gauge needle, squeezed gently to force out a small amount of feces, and then returned to the abdominal cavity. The abdominal incision was closed in layers, and the animals received 30 mL/kg body weight normal saline solution subcutaneously immediately after CLP as a fluid resuscitation. Two groups of rats with $n=12$ per group were used in this experiment. The treatment group received highly purified IaIp at 10 and 20 hrs after CLP at 30 mg/kg bodyweight. The control group received saline. At 20 hr after CLP, the necrotic caecum was excised and the abdominal cavity washed twice by using 40 mL of warm, sterile normal saline solution. The abdominal incision was then closed in layers. The procedure of cecal excision in CLP animals was performed to mimic the clinical situation in which the septic focus is removed. The experimental animals were allowed food ad libitum and monitored for 10 days to record the time of death for the non-survivors. A log-rank test was employed for comparison of mortality rates among different groups of animals and p values were determined by Kaplan-Meier method. A significant increase in survival was observed in the treatment group compared to the saline control group (81.3% of the animals in the treatment group survived vs. 44% in the control group (p value=0.0293)), suggesting that the highly purified IaIp is biologically active and effective in reducing the sepsis-related death in septic animals. Results are shown in Figure 9.

Table 1. Comparison of the specific inhibitory activity of IaIp

IaIp preparation Purified from	Protein conc. [mg/mL]	IaIp conc. [mg/mL]	IaIp purity [%]	Specific Inh. Activity [TIU/mg IaIp]
Cryoprecipitate	7.30	5.10	69.86%	1412 \pm 47.52
Cryopoor plasma	17.30	17.00	98.27%	1409 \pm 42.50

TIU=Trypsin inhibitory unit; mean \pm SD from three independent experiments.

Specific inhibitory activity of the highly purified IaIp (from cryo-poor plasma) was calculated and compared with IaIp purified from the cryoprecipitate, a side-fraction of fVIII production. The biological activity of IaIp was measured in a trypsin inhibition assay using the chromogenic substrate L-BAPA (N(alpha)-Benzoyl-L-arginine-4-nitroanilide hydrochloride (Fluka Chemicals). This assay is based on the ability of IaIp to inhibit the hydrolysis of L-BAPA. Inhibition can be monitored by a decrease in the rate of Δ absorbance/minute at 410 nm.

Total protein concentration was quantitatively measured by the BioRad protein assay and IαIp concentrations were measured by a competitive ELISA using MAb 69.31 as described in Lim *et al*, J. of Infectious Diseases, 2003). There was no significant difference in the specific trypsin inhibitory activity between IαIp preparations purified from cryo-poor plasma and cryoprecipitate
5 (*p* value= 0.939), suggesting that IαIp in both fractions had comparable biological activity.

EXAMPLE 7

Side fraction from FIX purification

The "washing fraction" from the chromatographic purification of clotting factor FIX on
10 DEAE-Sepharose FF (Josic et al. Journal of Chromatography, cited above). This fraction can be eluted with a 0.01 – 0.1 M sodium citrate/0.005-0.1 M sodium phosphate buffer, pH 6.0 containing 0.25 M sodium chloride. The IαI and PaI are the main components in this fraction. The FIX containing fraction can be eluted from DEAE Sepharose FF column with a 0.01-0.1 M sodium citrate/0.005-0.1 M sodium phosphate buffer, pH 6.0 containing 0.3-0.6 M sodium
15 chloride in the next step. This fraction also contains other vitamin K dependent clotting factors such as clotting factor II (FII), clotting factor X (FX), lower amounts of clotting factor VII (FVII) together with residual amounts of IαIp. After dialysis to reduce osmolality and salt concentration, the residual IαIp in the FIX fraction can be recovered by affinity chromatography on immobilized heparin and step elution in buffers with increasing salt concentration and
20 osmolality. Fractions from an early elution step in the wash buffer contain over 80% IαIp with very low FIX contamination. Elution of FIX occurs at a later step in a buffer with higher salt concentration and osmolality. There is also a flow through fraction that does not bind to heparin, is free of FIX and contains a mixture of IαIp (30-40%), vitamin K-dependent clotting factors, and the solvent/detergent (S/D) used for virus inactivation. The S/D can be removed in an
25 additional chromatographic step of DEAE-Sepharose FF.

IαIp containing fractions collected from DEAE-Sepharose, immobilized heparin or the flow through from heparin can be further purified individually or as a pool by hydroxylapatite chromatography. Using either approach, the final preparation contains more than 90% ITI.

The concentrate purified using these protocols contains more than 90% IaI/PaI. It has been virus inactivated by solvent/detergent treatment. Terminal heating in final container with or without use of stabilizers for more than 30 minutes or pasteurization at 55-65°C in the presence of stabilizers can be introduced as a second virus inactivation step without significant loss of activity.

The resulting concentrate containing more than 90% IaI/PaI can be virus inactivated with S/D treatment or as a second inactivation step, terminal heating of the purified proteins with or without stabilizers for 30 minutes or alternatively, pasteurization at 55-65 °C in the presence of stabilizers.

10 **A strong anion-exchange fraction**

Instead of the eluate after solid-phase extraction with weak anion-exchanger DEAE Sephadex A50,) an eluate after solid phase extraction with a strong anion-exchanger Q Sephadex A50 can be used. Conditions for elution are described in German patent DE 4342132C1. A mixture of IaI and PaI do not bind or only weakly binds to the monolithic anion-exchange support DEAE-CIM. Other proteins such as clotting factors FII, FVII, FIX and FX, clotting inhibitors PC, PS and PZ, adhesion protein vitronectin and protease FSAP are eluted in separate fractions. Further separations of the remaining contaminants can be achieved in the next step using hydroxylapatite chromatography.

A monolith chromatographic fraction

20 A DEAE CIM monolith (membrane) can be used for chromatographic separation in FIX purification instead of DEAE Sepharose FF or other particle-based anion exchanger (See DE 4342132C1). Surprisingly, IaI and PaI did not bind or bound only weakly to the monolithic support. Other proteins, such as FIX, vitronectin and FII, FVII (low amount) and FX were eluted as separate fractions. The proteolytic activity, coming mainly from Factor VII Activating
25 Protease (FSAP) (J. Roemisch, Biological Chemistry 383 (2002) 1119-1124) was also completely separated in this purification step. Further separation of the remaining contaminants from IaI/PaI containing fraction(s), namely traces of vitamin K dependant clotting factors FII, FVII and FX were achieved in the next step using hydroxylapatite chromatography.

All publications and patent documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication or patent document were so individually denoted. Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton *et al.*, *Dictionary of Microbiology and Molecular Biology* (2nd ed. 1994); *The Cambridge Dictionary of Science and Technology* (Walker ed., 1988); *The Glossary of Genetics*, 5th Ed., R. Rieger *et al.* (eds.), Springer Verlag (1991); and Hale & Marham, *The Harper Collins Dictionary of Biology* (1991).

INCORPORATION BY REFERENCE

The contents of all references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated herein in their entireties by reference.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.